



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

353. Proposed by RICHARD P. LOCHNER, Philadelphia, Pa.

The center of a sphere, radius $R = 5$ inches, is $a = 10$ inches above the surface of a sphere, radius $R = 12\frac{1}{2}$ inches. There is a point of light at $b = 1$ inch horizontally from a point $c = 10$ inches vertically above the surface of the first sphere. What is the area of the shadow which the upper sphere casts on the lower one?

354. Proposed by C. N. SCHMALL, New York City.

Prove $\Gamma(1+x)\Gamma(1-x) = \frac{\pi x}{\sin \pi x}$.

355. Proposed by C. N. SCHMALL, New York City.

Given the curve of the n th degree,

$$y^n - (a + bx)y^{n-1} + (c + dx + ex^2)y^{n-2} + \dots = 0,$$

show that if each ordinate is divided by the corresponding subtangent, the sum of all the resulting ratios will be a constant.

MECHANICS.

When this issue was made up no solutions had been received for numbers 271, 272, 275, 277, 278, 279, and 282.

286. Proposed by C. N. SCHMALL, New York City.

A slightly elastic string is just long enough to reach between two hooks on the same horizontal line. A ring of weight w is placed at its middle point. Show that the ring will sink through a distance $h = a\sqrt[3]{3ew/2}$, where e is the elasticity of the string and $2a$ the distance between the hooks.

287. Proposed by WALTER H. DRANE, Lebanon, Tenn.

While sitting in an empaled enclosure, I noticed that the spokes of the wheels of passing automobiles, when viewed through the pickets of the fence, appeared to revolve more slowly than they really did, and in some instances even appeared to be revolving in a direction opposite to that in which they were really turning. Explain this optical illusion.

NUMBER THEORY.

When this issue was made up no solutions had been received for numbers 187, 189, 191, 192, 194, 201, and 202 inclusive.

205. Proposed by E. T. BELL, New York City.

Show that in the usual arithmetical sense the form that follows admits of composition; give the requisite transformations, and indicate how several (if not all) solutions may be found. The variables are the x_i .

$$x_0^2 + nx_1^2 + mrx_2^2 + mnrx_3^2 + mnrx_4^2 + mn^2r^2x_5^2 + nr^2m^2x_6^2 + rm^2n^2x_7^2.$$

206. Proposed by R. D. CARMICHAEL, Indiana University.

Prove that the sum of the sixth powers of two integers cannot be the square of an integer.

207. Proposed by A. J. KEMPNER, University of Illinois.

There are 80 positive integers < 100 containing no figure 9 against 19 containing at least one figure 9. (For integers < 1000 the numbers are 728 and 271 respectively.) One might be led to believe that for every positive integer M the number N_1 of positive integers $< M$ containing no figure 9 is always greater than the number N_2 of positive integers $< M$ containing at least one figure 9.

To prove: $\lim_{M \rightarrow \infty} N_1/N_2 = 0$. See pages 48-50 of January, 1914, issue.